DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY

Prepared for the
NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

ATLAS OF MARS
1:15,000,000 GEOLOGIC SERIES
EASTERN REGION
M 15M, 0/270 G, 1987
I-1802-B

North

West

East

South

INDEX OF THE 1:15,000,000 GEOLOGIC SERIES

SCALE 1:15,000,000 (1 mm=15 km) AT 0° LATITUDE
MERCATOR PROJECTION

KILOMETERS

Prepared on behalf of the Planetary Geology and Geophysics Program, Planetary Division, Office of Space Science and Applications, National Aeronautics and Space Administration, under contract W-15,614 through Ames NCA2-1544 to Arizona State University.

CORRELATION OF MAP UNITS, GEOLOGIC EVENTS, AND CRATER DENSITIES IN THE EASTERN EQUATORIAL REGION OF MARS

NOTES ON BASE

This map sheet is one of a series covering the entire surface of Mars at a scale of 1:15,000,000. Sources for the map base were 1:5,000,000 scale shaded relief maps described by Batson and others (1979). Data used in the map portrayal were obtained from Viking Orbiter imagery.

ADOPTED FIGURE

The figure of Mars used for computing the map projection is an oblate spheroid (flattening of 1/192) with an equatorial radius of 3,393.4 km and a polar radius of 3,375.7 km.

PROJECTIONS

The Mercator projection is used between the 57° parallels; the Polar Stereographic projection is used for the polar regions between 57° and the poles. Scale is 1:15,000,000 at the equator and 1:9,000,000 at the 57° parallels. The projections have a common scale of 1:8,181,818 at lat ±54.09°. Longitude increases to the west in accordance with astronomical convention for Mars. Latitudes are areographic.

CONTROL

Planimetric control for the 1:15,000,000-scale maps used to compile the bases for these sheets was derived from photogrammetric triangulations by use of Mariner 9 pictures (Davies, 1973). This control network was densified through the use of Viking data (Davies and others, 1978). At least 80 percent of the image control points lie within 0.5 mm of the positions published in (1).

MAPPING TECHNIQUE

The mapping bases for this series were prepared by using computer-shaded relief maps reduced and digitally transformed where necessary to fit the projections used. Topographic data (shaded relief portrayal, features on these bases were used to position details taken from Viking Orbiter pictures. Features were drawn as if illuminated uniformly from the west through use of airbrush techniques described by Inge (1972) and photointerpretive methods described by Inge and Bridges (1976). The shading is not generalized and accurately represents the character of surface features.
Shaded relief analysis and portrayal were by Barbara J. Inge.

NOMENCLATURE

All names on this sheet are approved by the International Astronomical Union (IAU, 1974, 1977, 1980, 1985, and 1986). Named features and their positions are taken from published maps of Mars that have scales of 1:2,000,000, 1:5,000,000, and 1:25,000,000.
M 1204 0270 G

REFERENCES

Batson, R.M., Bridges, P.M., and Inge, J.L., 1979, Atlas of Mars, The 1:5,000,000 map series: National Aeronautics and Space Administration Special Publication 438, 146 p.
Davies, M.E., 1973, Mariner 9 mapping control net: Photogrammetric Engineering, v. 39, no. 12, p. 1287–1302.
Davies, M.E., Katayama, P.Y., and Roth, J.A., 1978, Control net of Mars: February 1978: The Rand Corporation, Publication R-2270-NASA, 27 p.
Inge, J.L., 1972, Principles of lunar illustration: Aeronautical Chart and Information Center Reference Publication 72-1, 22 p.
Inge, J.L., and Bridges, P.M., 1976, Applied photointerpretation for airbrush cartography: Photogrammetric Engineering and Remote Sensing, v. 42, no. 6, p. 789–796.
International Astronomical Union, 1974, Commission 16: Physical study of planets and satellites, and Lunar and martian nomenclature, in 15th General Assembly, Sydney, 1973, Proceedings: International Astronomical Union Transactions, v. 15B, p. 100–158, 271–321.
_____ 1977, Working Group for Planetary System Nomenclature, in 16th General Assembly, Grenoble, 1976, Proceedings: International Astronomical Union Transactions, v. 16B, p. 305–307.
_____ 1980, Working Group for Planetary System Nomenclature, in 17th General Assembly, Montreal, 1979, Proceedings: International Astronomical Union Transactions, v. 17B, p. 295–297.
_____ 1985, Working Group for Planetary System Nomenclature, in 18th General Assembly, Patras, 1982, Proceedings: International Astronomical Union Transactions, v. 18B, p. 314–317.
_____ 1986, Working Group for Planetary System Nomenclature, in 19th General Assembly, New Delhi, 1985, Proceedings: International Astronomical Union Transactions, v. 19B, p. 347–350.

INTRODUCTION

The Mariner 9 mission in the early 1970's provided the first comprehensive view of the geology of Mars (McCauley and others, 1972; Masursky, 1973) and led to the identification of the first global geologic map (Carr and others, 1973). These preliminary studies were followed by more comprehensive geologic mapping coordinated by the U.S. Geological Survey. Three quadrangles at a scale of 1:5,000,000 were produced in this map series, from which Scott and Carr (1978) compiled a single geologic map at a scale of 1:25,000,000. The reliability of these maps, however, is varied because of the uneven quality of the Mariner 9 data.

Viking Orbiter images fill important gaps in the Mariner data and provide substantially better coverage at both moderate (300 to 1,000 m/pixel) and high (better than 100 m/pixel) resolutions. The increased quality of the image results from a better camera system, improved image processing, and clearer atmospheric conditions than those of the Mariner 9 mission. These factors, along with the higher resolution obtained by the Viking Orbiter cameras, have enabled the discovery of geologic relations previously unknown. Photogeologic studies based on the Viking images, together with results from other remote-sensing experiments, such as the Infrared Thermal Mapper flown on Viking Orbiter (Kieffer and others, 1977), provide the basis for compilation (at 1:15,000,000 scale) of the three map sheets in the present series: the geologic map of the western equatorial region of Mars (Scott and Tanaka, 1986), the geologic maps of the polar regions of Mars (Tanaka and Scott, 1987), and this sheet.

Photogeologic mapping techniques employed here follow principles defined by Shoemaker and Hackman (1962) for the Moon and refined by Wilhelms (in press). Initial mapping was compiled on uncontrolled mosaics prepared during Viking mission operations. Although these mosaics were produced at a wide range of scales, were not uniform in format, and did not cover the entire area of interest, the preliminary mapping provided insight into local geologic relations. Results were then compiled on 1:2,000,000-scale photomosaics of moderate-resolution photographs (~300 to 100 m/pixel). The final ink compilation was prepared from the 1:2,000,000-scale photomosaics onto the shaded relief 1:15,000,000-scale map base (U.S. Geological Survey, 1982).

The map portrays inferred rock units, stratigraphic relations, and other features such as faults. From this mapping, the major impact, tectonic, and volcanic events that have shaped the surface of Mars are interpreted, as are modifications produced by wind, water, and other agents of geologic change. Definition of possible rock units is based on assessment of their geomorphology, albedo, spectral properties, and thermal characteristics. As discussed by Milton (1974), some units on Mars are characterized principally by their geomorphic expression, which locally may be due more to secondary modification than to their primary nature. Mapped units are assigned to time-stratigraphic systems formulated from Mariner 9 mapping (Scott and Carr, 1978), which were adapted for the map of the western equatorial region (Scott and Tanaka, 1986) and refined by Tanaka (1986).

Relative ages of the rock units are established by stratigraphic relations, cross-cutting relations of structural features, and size-frequency distributions of impact craters. Ages based on crater frequency must be employed with caution, however, especially in areas where wind and water have modified the surface and where sedimentary deposits may have partly buried other units. As described by Condit (1978), craters 4 to 10 km in diameter are important for use in broad relative-age dating of Martian surfaces; smaller craters may have been buried, eroded, or extensively modified, whereas larger craters are too scarce to produce statistically meaningful results. Crater-density numbers on the correlation chart were derived according to the convention established by Scott and Tanaka (1986). Because small craters are better preserved on the relatively smooth, younger geologic units than on the rough, uneven, or highly faulted older materials, different sizes of craters were counted for rock units in each system: greater than 2 km for the Amazonian, greater than 2 and 5 km for the Hesperian, and greater than 5 and 16 km for the Noachian.

Rock-stratigraphic classifications of diverse rocks (formations and members) are used to minimize cumbersome adjectival descriptions. The term "assemblage" (Scott and Tanaka, 1986) is employed for sets of units that share common attributes, for example, the Hellas assemblage encompasses those units associated with the formation and modification of the Hellas basin. Many of the units appear to intergrade, and some contacts are drawn primarily on the basis of secondary characteristics such as discontinuities in impact-crater density; these contacts are dashed.

Impact craters that are superposed on the surrounding terrain and have well-preserved, extensive rim units (referred to be ejecta deposits) are shown in yellow; only those craters having material units larger than 100 km across are mapped. Craters with rims that are partly buried by younger units and craters that are known in broken; only those craters having rim diameters larger than 150 km are mapped. Craters are not otherwise classified by type or by age.

PHYSIOGRAPHIC SETTING

Mars' surface can be divided into two areas that are distinguished by the Hesperian along the northern lowlands-southern highlands boundary and on the northeast margin of the Hellas basin. Channel formation along the boundary was concentrated near Deuteronilus Mensae and represents major modification of the scarp boundary. East of Deuteronilus, channels are not abundant; however, scarp modification or retreat is indicated by knobby material mapped as part of the undivided unit (unit HNu). Parts of the southern highlands have been modified by erosion to produce chaotic material (unit Hch).

The Tharsis region is characterized by the Amazonian and the northern lowlands as part of the northern plains assemblage. It consists of four materials representing various stages of degradation and modification: knobby member (unit Hvk), mottled member (unit Hvm), grooved member (unit Hvg), and ridged member (unit Hvr). The ridged material, the knobby member, and the mottled member are characterized by closely spaced, conical hills that appear darker than the surrounding terrain. The mottled member forms an extensive cover on the ridged material and the knobby member and has an albedo of 0.97 and 0.79, although Mariner 9 images indicate only a blurred surface having high albedo contrast; Viking images show that the contrasting albedo is due to impact craters whose ejecta deposits are brighter than intercrater areas (Hartmann, 1966, p. 296). The grooved member is characterized by irregular to polygonal patterns formed by grooves and troughs as long as 20 km, which may have been produced by peripheral, tectonic, or compaction processes (Carr and Schaber, 1977; Pechmann, 1980; McGill, 1985) enhanced by erosion. The ridged member is distinguished by wholer-like patterns of ridges resembling those in the Deuteronilus Mensae region. Stratigraphic relations between Vastitas Borealis Formation and the younger Arcadia Formation are unclear except in eastern Arcadia Planitia, where the Amazonian unit embays the ridged unit (Hvr). The Vastitas Borealis Formation may be locally younger than the Vastitas Borealis Formation.

Cratered terrains are not as obvious on Mars as on the Moon and Mercury. Although this may represent a difference in impact flux, a more likely explanation is that erosion and resurfacing have obliterated much of the early Martian crustal record. Fluvial processes are responsible for the remnants of possible basins other than Hellas and Isidis (such as Aram Chaos and the "Ladon basin") have been described by Wood and Head (1976) and Schultz and Glicken (1979).

STRATIGRAPHY AND HISTORY

Noachian System

The Noachian System comprises the oldest rocks discernible on Mars (Scott and Carr, 1978). Size frequencies of craters larger than 16 km in diameter on Noachian surfaces are similar to those of the Nectarian System on the Moon and may represent a similar period of impact cratering (Tanaka, 1986).

Two of the oldest rock units on Mars are basin-rim material (unit Nh1) and hilly material (unit Nplh), which form rugged, mountainous terrain associated with the Hellas and Isidis basins, respectively; a third ancient unit, mountainous material (unit Nm), forms massifs in the southern highlands. Because this unit does not everywhere appear to be associated with a clearly recognizable basin, but it is considered to represent parts of basins now grossly destroyed.

Overlying these rugged and intensely cratered terrains are old plateau units, including cratered material (unit Npc) and a subdued, cratered plateau material (unit Npl1). Both units are characterized by flood-style volcanism (Greeley and Spudis, 1981); such a volcanic origin is indicated by the presence of large mare-type (wrinkle) ridges in the ridged material (unit Npl2). The dissected material (unit Npld) has been cut by extensive channels that are presumably of fluvial origin; the etched material (unit Npla) appears to be marked by sinuous to braided patterns of fine ridges and troughs, in turn, these may represent exhumed and etched appearance.

Undivided material (unit HNu) consists of rock units in the highlands and lowlands that cannot be distinguished morphologically or placed precisely in a stratigraphic position in the Hesperian or Noachian Systems.

Rock units of Noachian age are considered to result from the final stages of planetary accretion and the waning period of heavy bombardment. In the middle to late Noachian Period, we infer the extrusion of extensive flood lavas and contemporaneous surface modifications by eolian, fluvial, and other surface processes that continued into the Hesperian System.

Hesperian System

As in the western hemisphere (Scott and Tanaka, 1986), many highland areas are interpreted to have been smoothed and resurfaced by lava flows and eolian deposits, representing a continuation of activity that began during the Noachian. Units of Hesperian age are distinguished by having 67 to 200 craters larger than 5 km in diameter per 10^6 km².

The most extensive plains unit in the Hesperian System is the *ridged plains material* (unit Hr), which has sinuous ridges resembling those on the lava flooded lunar maria and is inferred to be lava flows. It forms the basal unit of the Hesperian System (Scott and Carr, 1978). The largest concentration of ridged plains material in the map area is in Hesperia Planum northeast of the Hellas basin. In Syrtis Major Planum, a widespread occurrence of apparently younger material is designated the *Syrtis Major Formation* (unit Hs); unlike the ridged plains material (unit Hr), flow units of the Syrtis Major Formation have clearly distinguishable flow fronts and margins, and they appear to have originated from at least two central features resembling calderas (Schaber, 1982). Although these flows display many ridges, the ridges are generally narrower and shorter than those of the ridged plains northeast of Hellas.

In addition to eruptions in Syrtis Major, the middle Hesperian Period was marked by the initiation of central-vent volcanism in the Elysium region; lavas were extruded to form the *Hecates Tholus Formation* (unit Hhet) and the older flows of the *Albor Tholus Formation* (unit AHet). Eruptions also occurred in the southern hemisphere to form the dissected member of the *Amphitrites Formation* (unit Had) and most of the *Tyrrhena Patera Formation* (unit AHt), the *Apollinaris Patera Formation* (unit AHa), and the *Hadriaca Patera Formation* (unit AHh). These eruptions may be associated with crustal adjustments following the formation of the Hellas basin (Peterson, 1978). On the basis of photogeologic analysis, Greeley and Spudis (1981) postulated that Tyrrhena Patera was formed by magmas erupted through impact-generated regoliths that contained ground water. This process may have resulted in phreatomagmatic explosions that produced an early-stage ash shield, which was later degraded by surficial processes and which was the site of late-stage effusion of lavas. Although stages of Hesperian activity (at 67° S., long 299°), Apollinaris, and Hadriaca Paterae are inadequate for detailed analysis, a similar history is proposed for these eruptive centers.

Further resurfacing of the southern highlands is represented by the youngest units of the plateau sequence, which subdue or bury the ridged plains units. These units are characterized by smooth surfaces and are separated on the basis of albedo patterns into *smooth material* (unit Hplg) and *mottled plains material* (unit Hplm). Both units may consist of interbedded lava flows and sedimentary deposits; the mottling of the mottled plains unit may be caused by wind-sorted sediments.

The Hellas assemblage consists of rock units in and around Hellas Planitia. Its lowermost unit, the Hesperian basin-rim material noted above, is overlain by the *ridged plains floor unit* (unit Hfr), presumed to be flood lava-type counterparts of the ridged plains unit), which is in turn overlain by the *dissected floor unit* (unit Hhd), which may be of mixed origin.

Older channel material (unit Hch) was also emplaced during the Hesperian along the northern lowlands-southern highlands boundary and on the northeast margin of the Hellas basin. Channel formation along the boundary was concentrated near Deuteronilus Mensae and represents major modification of the scarp boundary. East of Deuteronilus, channels are not abundant; however, scarp modification or retreat is indicated by knobby material mapped as part of the undivided unit (unit HNu). Parts of the southern highlands have been modified by erosion to produce chaotic material (unit Hch).

The Tharsis region is characterized by the Amazonian and the northern lowlands as part of the northern plains assemblage. It consists of four materials representing various stages of degradation and modification: knobby member (unit Hvk), mottled member (unit Hvm), grooved member (unit Hvg), and ridged member (unit Hvr). The ridged material, the knobby member, and the mottled member are characterized by closely spaced, conical hills that appear darker than the surrounding terrain. The mottled member forms an extensive cover on the ridged material and the knobby member and has an albedo of 0.97 and 0.79, although Mariner 9 images indicate only a blurred surface having high albedo contrast; Viking images show that the contrasting albedo is due to impact craters whose ejecta deposits are brighter than intercrater areas (Hartmann, 1966, p. 296). The grooved member is characterized by irregular to polygonal patterns formed by grooves and troughs as long as 20 km, which may have been produced by peripheral, tectonic, or compaction processes (Carr and Schaber, 1977; Pechmann, 1980; McGill, 1985) enhanced by erosion. The ridged member is distinguished by wholer-like patterns of ridges resembling those in the Deuteronilus Mensae region. Stratigraphic relations between Vastitas Borealis Formation and the younger Arcadia Formation are unclear except in eastern Arcadia Planitia, where the Amazonian unit embays the ridged unit (Hvr). The Vastitas Borealis Formation may be locally younger than the Vastitas Borealis Formation.

Cornonocopiae units formation of the younger Vastitas Borealis members was the beginning of the development of the *etched plains material* (unit AHpe) in southern and western Elysium Planitia. This unit is characterized by irregular mesas and pits representing the erosion of a mantle, possibly by wind.

Amazonian System

This system was defined by Scott and Carr (1978) to include the youngest units on Mars, most of which occur in the northern lowlands. As noted on Mariner 9 images, the northern lowlands are partly covered, featureless plains. However, Viking data reveal more complex terrains and suggest diverse origins and modification by a variety of processes. Amazonian units are subdivided by age on the basis of frequency distributions of craters larger than 2 km in diameter.

The oldest unit mapped in the northern Amazonian System includes the Elysium Formation and materials formed during the modification of the Hellas basin. During the early part of the Amazonian, centralvent volcanism continued to construct Albor Tholus, Ita Tholus, as well as those of the Hecates Tholus Formation, are associated with the younger lava flows of member 1 (unit Ael1) of the Elysium Formation. These fresh-appearing flows are radial to the Elysium Mons edifice and are gradational with flows of member 2 (unit Ael2) that form the edifice; these younger flows also appear fresh and display channels and lavasses interpreted as collapsed lava tubes. Northwest of Elysium Mons, in Utopia Planitia, member 3 (unit Ael3) forms rugged plains having hummocky surfaces; the member appears to overlie and be gradational with member 1. The plains have been extensively modified by processes inferred to be fluvial, eolian, and periglacial. Part of this modification may be due to channels that also cut member 1. The channel material, mapped as member 4 (unit Ael4), is considered volcanic, possibly derived from lahars of Elysium Mons (Christiansen and Greeley, 1981).

Partial filling, mantling, and modification of the Hellas basin continued through the middle of the Amazonian Period to produce the younger units of the Hellas assemblage. Most of these units occur near the basin rim and are considered to be mantling material, some of which is eroded. The *lineated floor unit* (unit AHa) and the *channeled plains rim unit* (unit AHg) have similar crater-count ages and may represent only slightly modified areas of a mantle. The lineated floor unit is characterized by straight and curvilinear lineaments within smooth plains material and may have formed by tectonic modification of mantling material. The channeled plains rim unit has a subdued appearance and low remnant mesas and narrow channels. The other younger Hellas units form plains distinguished by low-relief features that may reflect differential erosion of mantling material. These units are *reticulate floor material* (unit AHg) characterized by a reticulate pattern of ridges, a *rugged floor unit* (unit AHy) whose undulating surface is rugged on a kilometer scale, and a *knobby plains floor unit* (unit AHg) that forms low knobs a few kilometers across.

In the northern plains assemblage, the *Arcadia Formation* is inferred to consist of lava flows whose age range defines and spans the Amazonian Period (Scott and Carr, 1978). Of the five members present in the western equatorial region (Scott and Tanaka, 1986), only three occur in the eastern region: member 2 (unit Aa2), member 3 (unit Aa3), and member 4 (unit Aa4). Lobate flow fronts, crescentic lava-flow ridges, and features presumed to be lava channels are visible on high-resolution images and are considered to be evidence of volcanic origin.

The Medusae Fossae Formation (Scott and Tanaka, 1986) consists of three members distributed along an east-west border zone between the highland plateaus and lowland plains of Amazonis Planitia in the eastern part of the map area. The lower, middle, and upper members (units Aml, Amm, and Amu, respectively) are distinguished on the basis of morphology, stratigraphic relations, and albedo. The materials are seen on high-resolution Viking images as massive horizontal sheets that may have a combined thickness in excess of 3 km. They have smooth and flat to gently rolling surfaces that are breached in places where an upper member has been stripped by wind. Eolian erosion has also produced yardangs on the edges of some high-standing deposits of the upper member, which suggests that they are composed of friable materials (Ward, 1979). The deposits are less hilly and cratered than those of the highlands, but they have higher relief and a higher albedo than the plains materials. Although Scott and Tanaka (1982) preferred a volcanic origin for this formation, suggesting that it consists of ignimbrites and lava flows, it may be formed of vast accumulations of eolian deposits transported and trapped along the highland-lowland boundary (Lee and others, 1982; Thomas, 1982) or of materials related to ancient polar deposits (Schultz and Lutz-Garihan, 1981).

Resurfacing or modification of parts of the northern lowlands continued through the Amazonian Period to produce plains units characterized by diverse origins. These materials have been divided on the basis of smooth plains unit (unit Aps) and a *knobby plains unit* (unit Apk). In many places the smooth plains unit probably consists of eolian deposits. The knobby plains unit appears to be formed by the erosion of older units; surfaces between the knobs may be erosional or may consist of volcanic, mass-wasting, or eolian materials.

Surficial materials of inferred mass-wasting, fluvial, and eolian origin occur locally throughout the northern lowlands. Although they consist of interbedded lava flows and sedimentary deposits, the resurfacing of the mottled plains unit may be caused by wind-sorted sediments.

The Hellas assemblage consists of rock units in and around Hellas Planitia. Its lowermost unit, the Hesperian basin-rim material noted above, is overlain by the *ridged plains floor unit* (unit Hfr), presumed to be flood lava-type counterparts of the ridged plains unit), which is in turn overlain by the *dissected floor unit* (unit Hhd), which may be of mixed origin.

Older channel material (unit Hch) was also emplaced during the Hesperian along the northern lowlands-southern highlands boundary and on the northeast margin of the Hellas basin. Channel formation along the boundary was concentrated near Deuteronilus Mensae and represents major modification of the scarp boundary. East of Deuteronilus, channels are not abundant; however, scarp modification or retreat is indicated by knobby material mapped as part of the undivided unit (unit HNu). Parts of the southern highlands have been modified by erosion to produce chaotic material (unit Hch).

The Tharsis region is characterized by the Amazonian and the northern lowlands as part of the northern plains assemblage. It consists of four materials representing various stages of degradation and modification: knobby member (unit Hvk), mottled member (unit Hvm), grooved member (unit Hvg), and ridged member (unit Hvr). The ridged material, the knobby member, and the mottled member are characterized by closely spaced, conical hills that appear darker than the surrounding terrain. The mottled member forms an extensive cover on the ridged material and the knobby member and has an albedo of 0.97 and 0.79, although Mariner 9 images indicate only a blurred surface having high albedo contrast; Viking images show that the contrasting albedo is due to impact craters whose ejecta deposits are brighter than intercrater areas (Hartmann, 1966, p. 296). The grooved member is characterized by irregular to polygonal patterns formed by grooves and troughs as long as 20 km, which may have been produced by peripheral, tectonic, or compaction processes (Carr and Schaber, 1977; Pechmann, 1980; McGill, 1985) enhanced by erosion. The ridged member is distinguished by wholer-like patterns of ridges resembling those in the Deuteronilus Mensae region. Stratigraphic relations between Vastitas Borealis Formation and the younger Arcadia Formation are unclear except in eastern Arcadia Planitia, where the Amazonian unit embays the ridged unit (Hvr). The Vastitas Borealis Formation may be locally younger than the Vastitas Borealis Formation.

Cornonocopiae units formation of the younger Vastitas Borealis members was the beginning of the development of the *etched plains material* (unit AHpe) in southern and western Elysium Planitia. This unit is characterized by irregular mesas and pits representing the erosion of a mantle, possibly by wind.

Amazonian System

This system was defined by Scott and Carr (1978) to include the youngest units on Mars, most of which occur in the northern lowlands. As noted on Mariner 9 images, the northern lowlands are partly covered, featureless plains. However, Viking data reveal more complex terrains and suggest diverse origins and modification by a variety of processes. Amazonian units are subdivided by age on the basis of frequency distributions of craters larger than 2 km in diameter.

The oldest unit mapped in the northern Amazonian System includes the Elysium Formation and materials formed during the modification of the Hellas basin. During the early part of the Amazonian, centralvent volcanism continued to construct Albor Tholus, Ita Tholus, as well as those of the Hecates Tholus Formation, are associated with the younger lava flows of member 1 (unit Ael1) of the Elysium Formation. These fresh-appearing flows are radial to the Elysium Mons edifice and are gradational with flows of member 2 (unit Ael2) that form the edifice; these younger flows also appear fresh and display channels and lavasses interpreted as collapsed lava tubes. Northwest of Elysium Mons, in Utopia Planitia, member 3 (unit Ael3) forms rugged plains having hummocky surfaces; the member appears to overlie and be gradational with member 1. The plains have been extensively modified by processes inferred to be fluvial, eolian, and periglacial. Part of this modification may be due to channels that also cut member 1. The channel material, mapped as member 4 (unit Ael4), is considered volcanic, possibly derived from lahars of Elysium Mons (Christiansen and Greeley, 1981).

Partial filling, mantling, and modification of the Hellas basin continued through the middle of the Amazonian Period to produce the younger units of the Hellas assemblage. Most of these units occur near the basin rim and are considered to be mantling material, some of which is eroded. The *lineated floor unit* (unit AHa) and the *channeled plains rim unit* (unit AHg) have similar crater-count ages and may represent only slightly modified areas of a mantle. The lineated floor unit is characterized by straight and curvilinear lineaments within smooth plains material and may have formed by tectonic modification of mantling material. The channeled plains rim unit has a subdued appearance and low remnant mesas and narrow channels. The other younger Hellas units form plains distinguished by low-relief features that may reflect differential erosion of mantling material. These units are *reticulate floor material* (unit AHg) characterized by a reticulate pattern of ridges, a *rugged floor unit* (unit AHy) whose undulating surface is rugged on a kilometer scale, and a *knobby plains floor unit* (unit AHg) that forms low knobs a few kilometers across.

In the northern plains assemblage, the *Arcadia Formation* is inferred to consist of lava flows whose age

DESCRIPTION OF MAP UNITS

Unit descriptions are based on characteristics observed primarily on Viking Orbiter images, supplemented by data from Mariner 9 and Earth-based telescope-sensing instruments such as the Viking Infrared Thermal Mapper. Most units are grouped by geologic assemblages, according to the convention of Scott and Tanaka (1986).

SURFICIAL DEPOSITS

As SLIDE MATERIAL—Smooth to lobate flow material associated with some scarps and crater rims; especially well developed along channel walls in Elysium Mons and Olympus Mons. Forms aprons along scarps and around knobs and mesas; may completely cover these features. Primarily equivalent to third type of slide material mapped by Scott and Tanaka (1986). No specific type area. *Interpretation:* Unconsolidated material, such as debris flows, resulting from mass wasting.

Asv YOUNGER CHANNEL AND FLOOD-PLAIN MATERIAL, DIVIDED—In eastern equatorial part of map area; forms plain as wide as 600 km marked by dark, sinuous, intertwining albedo patterns; appears more mottled westward. Type areas (Scott and Tanaka, 1986): lat 10° N., long 177° (younger channel material) and lat. 22° N., long 171° (flood-plain material). *Interpretation:* Fluvial deposits; distinct albedo patterns probably represent contrasts between dark and bland deposits; mottled cover in western part may represent deposition from ponded terminus of fluvial system.

Asl DUNE MATERIAL—Patches of barchan and other types of dunes on crater floors. Makes up part of dunes and dune-capped material (unit Ad) of south polar region (Tanaka and Scott, 1987). Type area: lat 48° S., long 109°. *Interpretation:* Sand deposited in areas of low wind strength.

LOWLAND TERRAIN MATERIALS

Consist of all plains-forming units between highland-lowland boundary scarp and north edge of map area, as well as materials of certain volcanic assemblages.

Northern plains assemblage

Materials deposited in widespread sheets on northern plains. Within each formation, members mapped on basis of morphology, albedo, and crater size-frequency distribution; some contacts approximately located. This assemblage underlies formation of highland-lowland boundary scarp (Scott, 1979).

Aps SMOOTH PLAINS MATERIAL—Forms patches and regions of flat, featureless plains; lightly cratered. Type areas: lat 9° N., long 222°. *Interpretation:* Probably of volcanic origin; fluvial processes probably consist of eolian deposits.

Apkpe ETCHED PLAINS MATERIAL—Occurs as patches in Elysium Planitia. Surface characterized by irregular mesas and pits. Type area: lat 15° N., long 238°. *Interpretation:* Plains deposits mantled by eolian material that has subsequently been eroded, possibly by wind.

Aa4 Aa3 Aa2 Aa1 ARCADIA FORMATION—Forms low-lying plains (1) in Arcadia Planitia east of Phlegra Montes in east margin of map area and (2) in Amazonis Planitia and parts of western map area. Common boundaries of older members mapped arbitrarily in places. Flows with lobate margins, ridges, and channels and small hills with summit craters which locally are breached and (or) etched by Scott and Tanaka (1986). No specific type area. Interpretation: Mostly lava flows and small volcanoes

Member 4—In Arcadia Planitia overlies member 3. Type area: lat 40° N., long 175° (Scott and Tanaka, 1986)

Member 3—In and north of Arcadia Planitia along east margin of map area. Flow fronts visible in places. Type area: lat 15° N., long 155° (Scott and Tanaka, 1986)

Member 2—Near type area. Type area: lat 30° N., long 40° (Scott and Tanaka, 1986)

Amu Amm Aml MEDUSAE FOSSAE FORMATION—Occurs near equator in eastern part of map area. Consists of extensive, relatively flat plains; may be composed in part of aeolian deposits. Subdivided on basis of albedo (Scott and Tanaka, 1986). Locally, where upper or middle member stripped by wind, underlying member shows inselbergs

Upper member—Surfaces smooth, flat to rolling, light in color; sculptured into ridges and grooves in places; broadly crater and mesa plains, locally mantled by younger units. Type area: lat 0° N., long 160° (Scott and Tanaka, 1986). *Interpretation:* Thick deposits of eolian sediments or volcanic pyroclastic deposits; wind eroded, partly mantled by younger volcanic rocks

Middle member—Similar to upper member but in places surface is more rugged and eroded. Type area: lat 10° N., long 160° (Scott and Tanaka, 1986). *Interpretation:* Forms by same processes as upper member, including indurated eolian or pyroclastic deposits; wind eroded, particularly along margins

Lower member—Most rugged unit; characterized by deeply eroded surfaces smooth to rough and highly eroded, darker than those of other members. One area: lat 3° S., long 182° contains long, broad troughs. Type area: lat 5° S., long 174° (Scott and Tanaka, 1986). *Interpretation:* Lava flows interbedded with eolian or pyroclastic deposits, in places heavily eroded

Aa VASTITAS BOREALIS FORMATION—Subpolar plains deposits of northern lowlands; members distinguished on basis of morphology and albedo contrast; placement of contacts locally arbitrary (Scott and Tanaka, 1986)

Hvm Mottled member—Major occurrence north of map boundary (Tanaka and Scott, 1987); extends as far south as about lat 40° N. Crosses Arcadia Planitia, where it embays the ridged material. Type area: lat 50° N., long 30°. *Interpretation:* Eolian deposits; in places may be underlain by volcanic rocks

Hvg Grooved member—Occurs as isolated patches in several areas of lowland plains; similar to mottled member but marked by curvilinear and polygonal patterns of grooves and troughs; closed polygons as wide as 20 km. Ridges present in some areas, as at lat 50° N., long 290°. Type area: lat 60° N., long 15° (Scott and Tanaka, 1986). *Interpretation:* Material same as mottled member; patterns may be due to compaction, tectonic, or volcanic processes

Hvr Ridged member—In isolated patches in northern plains. Characterized by concentric, low ridges about 1 to 2 km wide separated by intervening troughs. Commonly occurs as inliers in the lobate basin, mottled, and grooved materials; displays low mounds, many of which are aligned. Type area: lat 38° N., long 33° (Scott and Tanaka, 1986). *Interpretation:* Material same as mottled member; unit appears to develop from erosion of surrounding units. Origin of ridges unknown but they may result from peripiglacial or erosional processes. Low mounds in places may be remnants of volcanic cones or pyroclastic deposits

Hvk Knobby member—Similar in appearance to mottled member but generally has higher albedo and abundant small hills, some with summit craters. Crater ejecta have albedo similar to that of surrounding terrain. Type area: lat 55° N., long 5° (Scott and Tanaka, 1986). *Interpretation:* Plains of diverse origins (volcanic flows, eolian deposits); hills may be volcanic cones, remnants of highland terrain or of crater rims, or ejingas

Eastern volcanic assemblage

Volcanoes and lava flows in Elysium region (Greeley and Spudis, 1981).
ELYSIUM FORMATION—Lava flows associated with Elysium Mons and Albor Tholus.

Ael4 Member 4—Channel material. Type area: lat 36.9° N., long 209°. *Interpretation:* May be derived from lahars of members 1 and 3 (Christiansen and Greeley, 1981)

Ael3 Member 3—Forms plains having rugged relief, hummocky surfaces; lobate deposits seen at high resolution. Type area: lat 43° N., long 208°. *Interpretation:* Of volcanic origin; flows possibly derived from Elysium Mons, possibly interfinger with member 1. Extensively modified by fluvial, eolian, and periglacial processes

Ael2 Member 2—Lobate deposits, with rilles, comprising Elysium Mons edifice. Gradational with member 1. Type area: lat 25° N., long 215°. *Interpretation:* Lava flows displaying channels and partly collapsed lava tubes

Ael1 Member 1—Lobate, plains-forming deposits that radiate from Elysium Mons and overlie and embay Albor Tholus and Hecates Tholus Formations. Type area: lat 32.5° N., long 214°. *Interpretation:* Volcanic flows and related materials

AHet ALBOR THOLUS FORMATION—Material forming Albor Tholus; hummocky texture, more subdued than Hecates Tholus material. Type area: lat 18° N., long 210°. *Interpretation:* Volcanic flows

Hhet HECATES THOLUS FORMATION—Forms Hecates Tholus; hummocky surface cut by many narrow, sinuous channels. Type area: lat 30° N., long 209°. *Interpretation:* Volcanic flows, some emplaced through lava channels

Hs SYRTI