


Mercaator projection
Geodetic Reference System 1980, North American Datum 1983
Longitude of central meridian 70°39' W; latitude of true scale 41°39' N.
False easting 0 m; false northing 0 m
This map is not intended for navigational purposes.



DISCUSSION

Introduction

This map shows the sea floor in shaded relief view at a scale of 1:25,000, with sea floor depth as topographic contours overlain in blue. It is based on multibeam echo-sounder data collected during four cruises conducted between the fall of 1994 and the fall of 1998. The map is part of a 3-quadrangle map series showing the area offshore of Boston, Mass., that is comparable to the Stellwagen Bank National Marine Sanctuary map series (Valentine and others, 2001, 2003a-c; also see location map). Other maps of Quadrangle 2 depict topographic contours (Butman and others, 2003a), and backscatter intensity over shaded relief and topographic contours (Butman and others, 2003b).
The multibeam echo-sounder surveys were conducted aboard the vessel Frederick G. Creed, a SWATH Small Waterplane Area Twin Hull ship that surveys at speeds up to 15 knots. A Simrad Subsea EM 1000 Multibeam Echo Sounder (ES HFI), mounted on the starboard portion of the Creed, was used to acquire these data. The bathymetric soundings were gridded at 6 m (pixel resolution and smoothed using a 5-cell by 5-cell median filter; contours having a 5-meter interval were generated from the resulting grid). The shaded relief image was created by vertically exaggerating the topography four times and then artificially illuminating the relief by a light source positioned 45 degrees above the horizon from an azimuth of 350 degrees. In the resulting image, topographic features are enhanced by strong illumination on the northeast-facing slopes and by shadows cast on southwest-facing slopes. The image accentuates small features that could not be effectively shown by contours alone at this scale. Blank areas in the image represent places where no data exist. The two narrow strips of data in the southwest corner of the quadrangle were collected along single ship transits to Boston Harbor. Topographic lows are identified by backcast contours (backcasts face deeper water).

Some features in the image are artifacts of data collection and environmental conditions. They include small highs and lows and unnatural-looking features, and patterns oriented parallel or perpendicular to survey tracklines tracklines were run north-south in the eastern and north-central part of the map area, northwest-southeast in the south-central part, and northeast-southwest in the western part). For example, the winkle-like features about 100 m long, oriented east-west and perpendicular to the ship's track between 70°39' W and 70°42' W, are a result of heave of the vessel during data collection caused by large surface waves; the northeast-southwest-trending lines in the vicinity of 42°21.5' N, 70°47.5' W, are a result of data loss in the far range of individual swathes; and small features in features in the southwestern part of the study area, most notable along features that have sharp transitions, are a result of errors in synchronization of the navigation and multibeam

clinks during the 1994 survey (a diamond-shaped area with corners at 42°19.5' N, 70°48' W; 42°23' N, 70°45.5' W; 42°26' N, 70°48' W; and 42°23' N, 70°52.5' W). For example, see the northeast-southwest offsets of the northwest-southeast trending features near 42°21.94' N, 70°47.88' W, and near 42°23.74' N, 70°48.59' W.

Geologic history

The major topographic features shown in this map series were formed by glacial processes that occurred in several stages. Ice containing rock debris moved across the region, sculpting its surface and depositing sediment, forming the ridges and valleys that characterize the region. Other features are the result of processes at work when much of the area was covered with rotating ice, and when at the same time small valley glaciers and ice falls were active. Ice retreat and marine submergence occurred between 15 and 14 ka, resulting in a highstand of sea level approximately 33 m above modern sea level about 14 ka (Oldale and others, 1993). A lowstand of sea level approximately 45 m below modern sea level occurred about 12 ka as the earth's crust rebounded from ice loading. Thus, the sea floor of Massachusetts Bay in water depths shallower than about 45 m was reworked during the marine transgression between 18 and 14 ka, again during a relatively rapid sea-level regression between 14 and 12 ka, and finally during the transgression between 12 ka and the present. Today, the surficial sediments and features are reworked and shaped by tidal and storm-generated currents, which erode and transport sediments from the shallow areas into the deeper basins. Over time, the shallow areas affected by these processes were coarsened as sand and mud are removed and gravel remains, and the deeper basins have been built up as they receive the sand and mud. Knobel and Crow (1995) have identified areas of erosion, sediment reworking, and deposition in this region.

Quadrangle 2 features

One of the most striking aspects of the sea floor shown by this survey is the variability in bottom morphology and texture over scales of a few kilometers or less, caused by both natural and anthropogenic processes. The topography, surface features, and surficial sediment textures are the result of glacial processes, reworking during the last rise in sea level, reworking by modern processes, and the disposal of dredged and other material in this region over the last century.
West of about 70°40' W, the sea floor is characterized by a varying topography having relief of 10 to 15 m. North of about 42°23' N, a series of northwest-southeast-trending features that are approximately 2 km long and 0.5 km wide typically rise 5 to 10 m above the surrounding sea floor. Oldale and others (1994) examined 11 of these features in the region centered at 42°25' N, 70°48' W, originally mapped by Bothner and others, 1992, and hypothesized that they are submerged drumlins. In general, their tops are smooth (that is, little texture is evident

in the shaded relief image) in water deeper than 45 m, and rough in water shallower than 45 m. It is hypothesized that the rougher surface is the result of reworking and removal of fine material as sea level transgressed the region from its ~45 m lowstand.

South of 42°22' N and west of 70°41' W, the sea floor is dominated by a discontinuous and in places poorly defined ridge that extends roughly east-west. Its depth ranges from 40 to 45 m in the east to about 25 m in the west. Eastward of 70°41' W, the sea floor gradually deepens toward Stellwagen Basin. The sea floor is relatively flat and featureless with the exception of a thin, northwest-southeast-trending feature centered near 42°22.8' N, 70°38.15' W. This ridge, having relief of about 5 m, is steeper and narrower than the ridges to the west. In addition, a knob approximately 1 km in diameter and centered near 42°21.5' N, 70°36.5' W, rises about 10 m above the surrounding sea floor.

Several areas are hypothesized to be the remnants of meandering channels, which possibly are drainage routes formed during a lower stand of sea level. One channel starts at about 40 m water depth near 42°25.4' N, 70°45.8' W, and ends at about 50 m water depth in a lobe-like feature centered at about 42°25.4' N, 70°42.6' W. The channel is 100 to 200 m wide. A second, narrower channel starts near 42°22.2' N, 70°45.0' W, at about 35 m water depth, and ends near 42°22.15' N, 70°43.1' W, in water depth between 45 and 50 m.

Several areas suggest active sediment movement. Ribbon-like features, typically 0.5 to 2 km long and less than 40 m wide, are observed near 42°19.8' N, 70°46.9' W, and are suggestive of downslope sediment transport. The relief across these features is about 30 cm or less. Similar features are observed in the vicinity of 42°25.0' N, 70°46.5' W. Both sets of features trend generally northeast-southwest.

Four areas of the sea floor in Quadrangle 2 show the effects of ocean disposal of anthropogenic material. These areas are typically identified by a low mound having an unnatural-appearing roughness resulting from numerous individual dumps of material. The areas impacted by disposal are (1) centered near 42°26.0' N, 70°48.0' W; (2) located between 42°24.5' N, 70°49.0' W, and 42°25.5' N, 70°50.0' W; (3) located between 42°21.0' N, 70°39.75' W, and 42°21.5' N, 70°40.5' W; and (4) located between 42°19.5' N, 70°45.0' W, and 42°20.25' N, 70°46.0' W, at sites 1, 2, and 6, and the accumulation of material is a few meters higher than the surrounding sea floor. In the backscatter intensity image (Butman and others, 2003 b), numerous individual dumps are clearly visible as high-backscatter features. Areas 1 and 3 are within decommissioned dumping grounds, as shown on National Ocean Service (1997) Chart 13267, and area 2 is near the location of a former dumping ground barge (Butman and Lindsay, 1999). Area 4 is located to the south of the previous location of the Boston Lightship. Numerous additional isolated dumps are observed in a corridor between areas 3 and 4. A fifth area affected by disposal, observable in the backscatter intensity image but not in this shaded relief image, is located in the

northeastern part of the map area between 42°24' N, 70°36' W and 42°26' N, 70°39' W (see Butman and others, 2003b; Valentine and others, 1996).

The 50 individual diffuser heads for the new ocean outfall that discharges treated sewage effluent from the Boston metropolitan region into Massachusetts Bay extend between 42°23.06' N, 70°48.23' W, and 42°23.33' N, 70°46.81' W. The heads are located in a 30 to 35 m water depth in a topographic low, west-northwest-trending ridges rise to about 25 m below the sea surface to the north and south. The most notable features are two parallel rows, about 2 km long, of individual mounds, which consist of material discarded on the sea floor from the holes drilled for the risers that extend to the outfall tunnel below. The diffuser heads, which are about 3 m high and 4 m in diameter, are located between the rows and are not well resolved in the 6-meter-gridded data. Note that a single survey trackline was run over the diffuser heads' location in 1997. This data replaced the original survey in this area, eliminating the offsets caused by errors in the synchronization of the navigation and multibeam clocks during the 1994 survey.

ACKNOWLEDGMENTS

The multibeam echo-sounder surveys were conducted with support from the Canadian Hydrographic Service and the University of New Brunswick. For their skillful work at sea, we thank G. Costello, N. Doucet, J. Gagne, R. Sanficon, and B. Tester of the Canadian Hydrographic Service, E. Bedford of the National Oceanic and Atmospheric Administration, and the officers and crew of the Canadian Hydrographic Service survey vessel Frederick G. Creed. We also thank J.H. Clark of the University of New Brunswick for his assistance in data acquisition, processing, and interpretation.

REFERENCES CITED

Bothner, M.H., Parmenter, C.M., Twichell, D.C., Polton, C.F., and Knobel, H.J., 1992, A geologic map of the sea floor in western Massachusetts Bay, constructed from digital sidescan sonar images, photography, and sediment samples. U.S. Geological Survey Digital Data Series DDS-3, 1 CD-ROM.
Butman, Bradford, Hayes, Laura, Danforth, W.W., and Valentine, P.C., 2003a, Sea floor topography of Quadrangle 2 in western Massachusetts Bay offshore of Boston, Massachusetts. U.S. Geological Survey Geologic Investigations Series Map I-2732-A, scale 1:25,000.
—, 2003b, Backscatter intensity shaded relief, and sea floor topography of Quadrangle 2 in western Massachusetts Bay offshore of Boston, Massachusetts. U.S. Geological Survey Geologic Investigations Series Map I-2732-C, scale 1:25,000.

—, 2003c, Shaded relief and sea floor topography of Quadrangle 1 in western Massachusetts Bay offshore of Boston, Massachusetts. U.S. Geological Survey Geologic Investigations Series Map I-2731-B, scale 1:25,000.

Butman, Bradford, and Lindsay, John, eds., 1999, A marine GIS library for Massachusetts Bay focusing on disposal sites, contaminated sediments, and sea floor mapping. U.S. Geological Survey Open-File Report 99-439, 1 CD-ROM.
Knobel, H.J., and Crow, R.C., 1995, Seafloor environments within Boston Harbor—Massachusetts Bay sedimentary system, a regional synthesis. *Journal of Coastal Research*, v. 11, no. 1, p. 230-251.


National Ocean Service, 1997, Massachusetts Bay. National Oceanic and Atmospheric Administration, National Ocean Service, Chart 13267, scale 1:80,000.
Oldale, R.N., Colman, S.M., and Jones, G.A., 1993, Radiocarbon ages from two submerged strandline features in the western Gulf of Maine and a sea level curve for the northeastern Massachusetts coastal region. *Quaternary Research*, v. 40, no. 1, p. 38-45.

Oldale, R.N., Knobel, H.J., and Bothner, M.H., 1994, Submerged and eroded drumlins off northeastern Massachusetts. *Geomorphology*, v. 9, no. 4, p. 301-309.

Valentine, P.C., Danforth, W.W., Roworth, E.T., and Stillman, S.T., 1996, Maps showing topography, backscatter, and interpretation of surficial features in the Massachusetts Bay Disposal Site region off Boston, Massachusetts. U.S. Geological Survey Open-File Report 96-273, scale 1:10,000 and 1:12,500, 2 sheets.

Valentine, P.C., Middleton, T.J., and Fuller, S.J., 2001, Maps showing sea floor topography, sun-illuminated sea floor topography, and backscatter intensity of the Stellwagen Bank National Marine Sanctuary region off Boston, Massachusetts. U.S. Geological Survey Open-File Report 00-410, 1 CD-ROM, scale 1:60,000.
Valentine, P.C., Baker, J.L., and Unger, T.S., 2003a, Sea floor topography of the Stellwagen Bank National Marine Sanctuary off Boston, Massachusetts. U.S. Geological Survey Geologic Investigations Series Map I-2676-A, scale 1:60,000.
Valentine, P.C., Unger, T.S., and Baker, J.L., 2003b, Sun-illuminated sea floor topography of the Stellwagen Bank National Marine Sanctuary off Boston, Massachusetts. U.S. Geological Survey Geologic Investigations Series Map I-2676-B, scale 1:60,000.

—, 2003c, Backscatter intensity and sun-illuminated sea floor topography of the Stellwagen Bank National Marine Sanctuary off Boston, Massachusetts. U.S. Geological Survey Geologic Investigations Series Map I-2676-C, scale 1:60,000.



Location map showing mapped Quadrangle 2 shaded in blue. The western Massachusetts Bay map series comprises quadrangles 1-3 (outlined in black); the maps for Quadrangle 2 include this map and Butman and others (2003a,b). The shaded relief and sea floor topography of Quadrangle 1 are shown in Butman and others (2003a). Quadrangles 1-18 (outlined in gray) compose the computerized Stellwagen Bank National Marine Sanctuary (SBNMS) map series. The shaded relief and sea floor topography of the entire area of quadrangles 1-18 is shown at scale 1:60,000 in Valentine and others (2001, 2003a); it is also shown by quadrangle at scale 1:25,000 as U.S. Geological Survey Geologic Investigations Series Maps I-2701 (for quadrangle 1) through I-2718 (for quadrangle 18). The SBNMS boundary is shown as a dashed line. Selected bathymetric contours are labeled in meters.

**SHADED RELIEF AND SEA FLOOR TOPOGRAPHY OF QUADRANGLE 2
IN WESTERN MASSACHUSETTS BAY OFFSHORE OF BOSTON, MASSACHUSETTS**

By

Bradford Butman, Laura Hayes, William W. Danforth, and Page C. Valentine

2003